Patient Information Sheet – Shake/Jarratt's Lotion

This medicine was compounded specifically for you in a pharmacy to fill the prescription your doctor wrote for you. It was specially made to meet your individual needs. For this reason, no standardized information or literature is available with your prescription. If you have not done so, please discuss this medicine with your doctor to ensure that you understand (1) why you have been prescribed a compounded medicine, (2) how to properly take this medicine, and (3) the interactions, if any, this medicine may have with any other medicines you are taking.

Compounding is a long-standing pharmacy practice that allows doctors to treat their patients' individual needs without being restricted only to off-the-shelf medicines or devices. This medicine was prepared in a compounding pharmacy to meet the specifications ordered by your doctor.

Call your doctor if you experience any side effects, if you are taking additional medicines that may interact with this compounded medicine, or if you have allergies or other medical conditions that should be noted. Call our pharmacists if information on the label is not clear to you or you have any concerns regarding precautions, ingredients, or proper storage. Our pharmacists are available to address any additional questions or concerns.

Medication: Shake/Jarratt's lotion contains the active ingredients nystatin (anti-fungal), polymyxin (anti-bacterial), bacitracin (anti-bacterial) and hydrocortisone (anti-inflammatory).

Use: Shake/Jarratt's lotion is used to treat the inflammation/infection associated with diaper rash.

Side Effects: Often, topical medications provide more localized effects, reducing systemic side effects by circumventing vital organs. An allergic reaction to this medication is unlikely, but seek immediate medical attention if your child experiences a rash, itching, swelling, dizziness, or trouble breathing.

Administration: This medication may be applied after a diaper change up to four times daily to localized skin inflammation which can include redness, swelling and/or itching.

Storage, Beyond Use Dating and Disposal: Shake/Jarratt's lotion should be stored at controlled room temperature (68˚-77˚F). The "Beyond Use Date" is the expiration date for compounded medicines. This topical preparation is stable for six months after the dispensing date, however, it is always best to refer to your medication's label for specific storage requirements and beyond use dating. Search "How to Dispose of Unused Medicines" at www.fda.gov for info on medication destruction.

NORTH LAMAR
4018 N. Lamar Blvd.
Austin, TX 78756
512.459.9090

SOUTH LAMAR
3801B S. Lamar Blvd.
Austin, TX 78704
512.444.8866

WESTLAKE
2805 Bee Caves Rd., Ste. 416
Austin, TX 78746
512.327.8877

LAKELINE
13860 Research Blvd., Unit C
Austin, TX 78717
512.219.9499

COMPOUNDING CENTER
4120 Commercial Center Dr. #400
Austin, TX 78744
512.646.4007

www.PeoplesRx.com

Disclaimer: This Patient Information Sheet explains the proper use and storage of the described medication and is provided as a service to our customers. This guideline is a summary of information and is not intended to replace or act as an official drug monograph. Patients with further questions or concerns specific to this compounded medication should consult their pharmacist. Patients with questions regarding therapy not discussed above should consult their doctor.

Last update: May 2014